

SKEDULERINGSSTATUS S0

EIENDOMSNAAM EN DOSEERVORM

Ultimag Tablette

SAMESTELLING

Filmbedekte tablette bevattende:

Magnesiumchloried B. P. 660 mg

Sinkoksied 6 mg

FARMAKOLOGIESE KLASSIFIKASIE

A 24 Aanvullende mineraalpreparate, elektroliete.

INDIKASIES

Vir die aanvulling van magnesium en sink waar onvoldoende inname voorkom.

KONTRA-INDIKASIES

Nierontoereikendheid, ontwatering en maag-letsels en -stoornisse.

Sensitiwiteit vir enige van die bestanddele.

DOSIS EN GEBRUIKSAANWYSINGS

Neem 2 tot 4 tablette daagliks. Daar word aanbeveel dat die hoofdosering saans geneem word.

NEWE-EFFEKTE EN SPESIALE VOORSORGMAATREËLS

Diarree, naarheid en dermongemak kan voorkom. Magnesiumchloried moet met voorsorg aan pasiënte wat digitalisglikosiede ontvang, toegedien word. Gelyktydige toediening met 'n sinkoutsout kan die effek van penisillamien laat afneem. Tetrasikliene kan 'n chelaatkompleks vorm met sink.

BEKENDE SIMPTOME VAN OORDOSERING EN BESONDERHEDE VAN DIE BEHANDELING DAARVAN

Simptome van hipermagnesemie kan die volgende insluit: blosing van die vel, dors, hipotensie a.g.v. perifere vasodilatasie, lomerigheid, verwarring, verlies van tendonrefleks a.g.v. neuromuskulêre blok, spiertamheid, asemhalingsonderdrukking, hartaritmieë, koma en hartstilstand. Behandeling is simptomeaties en ondersteunend.

IDENTIFIKASIE

Wit, kapsule-vormige tablette.

AANBIEDING

ULTIMAG TABLETTE word voorsien in bruin glasbottels bevattend 30, 60 en 100 tablette.

BERGINGSAAWYSINGS

Berg benede 25 °C op 'n koel droë plek.

HOU BUITE BEREIK VAN KINDERS.

REGISTRASIENOMMER

X/24/79

NAAM EN BESIGHEIDSADRES VAN DIE APPLIKANT

Mirren (Edms) Bpk

Goldenrylaan 18

Morehill

Benoni 1501

Suid-Afrika

DATUM VAN PUBLIKASIE VAN HIERDIE VOUBILJET

9 November 1989

UGT-PI-PR-15/01

Pro-Print

SCHEDULING STATUS S0

PROPRIETARY NAME AND DOSAGE FORM

Ultimag Tablets

COMPOSITION

Film coated tablets containing:

Magnesium chloride B. P.	660 mg
Zinc oxide	6 mg

PHARMACOLOGICAL CLASSIFICATION

A 24 Mineral substitutes and electrolytes.

INDICATIONS

For the supplementation of magnesium and zinc in cases of insufficient intake.

CONTRA-INDICATIONS

Renal impairment; dehydration and gastric lesions and disorders.
Sensitivity to any of the ingredients.

DOSAGE AND DIRECTIONS FOR USE

Take 2 to 4 tablets daily. It is recommended to take the major dose at night.

SIDE EFFECTS AND SPECIAL PRECAUTIONS

Diarrhoea, nausea and bowel discomfort may occur. Magnesium chloride should be given with caution to patients receiving digitalis glycosides. Concurrent administration with zinc salt might diminish the effect of penicillamine. Tetracyclines may form a chelate with zinc.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT

Symptoms of hypermagnesaemia may include flushing of the skin, thirst, hypotension due to peripheral vasodilatation, drowsiness, confusion, loss of tendon reflexes due to neuromuscular blockade, muscle weakness, respiratory depression, cardiac arrhythmias, coma and cardiac arrest. Treatment is symptomatic and supportive.

IDENTIFICATION

White, capsule-form tablets.

PRESENTATION

ULTIMAG TABLETS are supplied in amber glass bottles containing 30, 60 and 100 tablets.

STORAGE INSTRUCTIONS

Store under 25 °C in a cool dry place.
KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER

X/24/79

NAME AND BUSINESS ADDRESS OF THE APPLICANT

Mirren (Pty) Ltd
18 Golden Drive
Morehill
Benoni 1501
South Africa

DATE OF PUBLICATION OF THIS PACKAGE INSERT

9 November 1989

UGT-PI-PR-15/01